

**Bogalecka Magda**

**Kołowrocki Krzysztof**

*Maritime University, Gdynia, Poland*

## **General model of critical infrastructure accident consequences application to chemical spill consequences generated by dynamic ship critical infrastructure network operating at the Baltic Sea waters. Part 3. Process of environment degradation**

### **Keywords**

Baltic Sea region, critical infrastructure, sea accident, accident consequences, environment degradation

### **Abstract**

In the paper, the process of environment degradation at the Baltic Sea area identification is performed. Next, the main characteristics of this process are predicted.

### **1. Introduction**

The probabilistic General Model of Critical Infrastructure Accident Consequences (GMCIAC) [Bogalecka, Kołowrocki, 2016], [Bogalecka, Kołowrocki, 2017c] includes the process of initiating events [Bogalecka, 2010], [Bogalecka, Kołowrocki, 2015a], the process of environment threats and the process of environment degradation [Bogalecka, Kołowrocki, 2015b] models.

The modelling critical infrastructure accident consequences was done in [EU-CIRCLE Report D3.3-GMU21, 2016] through designing the GMCIAC and the identification of its unknown parameters was performed in [EU-CIRCLE Report D3.3-GMU22, 2016]. Further, the GMCIAC adaptation to the prediction of critical infrastructure accident consequences was done in [EU-CIRCLE Report D3.3-GMU23, 2016].

### **2. Application of the model of the process of environment degradation to the Baltic Sea waters**

We assume, as in [EU-CIRCLE Report D3.3-GMU21, 2016], that the process of environment degradation of the sub-region  $D_k$ ,  $k = 1, 2, \dots, n_3$ , is taking  $\ell_k$ ,  $\ell_k \in N$ , different environment degradation

states  $r_{(k)}^1, r_{(k)}^2, \dots, r_{(k)}^{\ell_k}$ . Next, we mark by  $R_{(k/v)}(t)$ ,  $t \in (0, +\infty)$ ,  $k = 1, 2, \dots, n_3$ ,  $v = 1, 2, \dots, v_k$ , the conditional sub-process of the environment degradation of the sub-region  $D_k$ ,  $k = 1, 2, \dots, n_3$ , while the process of environment threats  $S_{(k)}(t)$  of the sub-region  $D_k$  is at the state  $s_{(k)}^v$ ,  $v = 1, 2, \dots, v_k$ .

The conditional sub-process  $R_{(k/v)}(t)$ , is a function defined on the time interval  $t \in (0, +\infty)$ , depending on the states of the process of environment threats  $S_{(k)}(t)$  and taking discrete values in the set  $\{r_{(k/v)}^1, r_{(k/v)}^2, \dots, r_{(k/v)}^{\ell_k}\}$  of the environment degradation states. We assume a semi-Markov model [Grabski, 2015], [Kołowrocki, 2004], [Kołowrocki, 2014], [Kołowrocki, Soszyńska-Budny, 2008], [Kołowrocki, Soszyńska-Budny, 2011], [Limnios, Oprisan, 2005], [Macci, 2008], [Mercier, 2008] of the sub-process of environment degradation  $R_{(k/v)}(t)$ , and we mark by  $\zeta_{(k/v)}^{ij}$  its random conditional sojourn times at the states  $r_{(k/v)}^i$ , when its next state is  $r_{(k/v)}^j$ ,  $i, j = 1, 2, \dots, \ell_k$ ,  $i \neq j$ ,  $k = 1, 2, \dots, n_3$ ,  $v = 1, 2, \dots, v_k$ . Under these assumption, the sub-process of environment degradation  $R_{(k/v)}(t)$ , for each sub-region  $D_k$ ,  $k = 1, 2, \dots, n_3$ , may be described by the vector

$[q_{(k/v)}(0)]_{1 \times \ell_k}$  of initial probabilities of the sub-process of environment degradation staying at particular environment degradation states at the initial moment  $t = 0$ , the matrix  $[q_{(k/v)}^{ij}]_{\ell_k \times \ell_k}$  of probabilities of transitions between the environment degradation states  $r_{(k/v)}^i$ , and  $r_{(k/v)}^j$ , and the matrix  $[G_{(k/v)}^{ij}(t)]_{\ell_k \times \ell_k}$ , of the distribution functions of the conditional sojourn times  $\zeta_{(k/v)}^{ij}$ , of the process  $R_{(k/v)}(t)$ , at the environment degradation states or equivalently by the matrix  $[g_{(k/v)}^{ij}(t)]_{\ell_k \times \ell_k}$ , of the density functions of the conditional sojourn times  $\zeta_{(k/v)}^{ij}$ ,  $i, j = 1, 2, \dots, \ell_k$ ,  $i \neq j$ ,  $k = 1, 2, \dots, n_3$ ,  $v = 1, 2, \dots, v_k$ , of the sub-process of environment degradation at the environment degradation states.

## 2.1. Parameters evaluation of the process of environment degradation at the Baltic Sea waters

To identify the unknown parameters of the process of environment degradation the suitable statistical data coming from realization should be collected. The statistical identification of the environment degradation was performed on the base on the ship accidents around the Baltic Sea in a period of 11 years (2004-2014). The initial moment  $t = 0$  of the process of environment degradation was fixed at the moment when the threat caused by ship accident generated one of the distinguished degradation effects states. Unfortunately, the less accurate identification of the process of environment degradation is performed for the Baltic Sea waters because of the less sufficiently numerous set of statistical data.

### 2.1.1. States of the process of environment degradation

Taking into account the expert opinion on varying in time the process of environment degradation, we distinguished its states for particular sub-regions. There are  $\ell_1 = 30$  states of the process of environment degradation in the air ( $D_1$  sub-region):

state  $r_{(1)}^1$  – accident has happened but it does not caused environment degradations,  
state  $r_{(1)}^2$  – the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(0, LC_{50}/2>$ ,

state  $r_{(1)}^3$  – the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}/2, LC_{50}>$ ,

state  $r_{(1)}^4$  – the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}, \infty)$ ,

state  $r_{(1)}^5$  – the realised substance caused the aesthetic nuisance of air of the accident area but the closure of area is not required,

state  $r_{(1)}^6$  – the realised substance caused the aesthetic nuisance of air of the accident area but the closure of area is not required, and simultaneously the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(0, LC_{50}/2>$ ,

state  $r_{(1)}^7$  – the realised substance caused the aesthetic nuisance of air of the accident area but the closure of area is not required and simultaneously the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}/2, LC_{50}>$ ,

state  $r_{(1)}^8$  – the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for not more than 2 days, and simultaneously the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}/2, LC_{50}>$ ,

state  $r_{(1)}^9$  – the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for not more than 2 days and simultaneously the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}, \infty)$ ,

state  $r_{(1)}^{10}$  – the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for 2 days or more, and simultaneously the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}, \infty)$ ,

state  $r_{(1)}^{11}$  – the realised substance changed pH regime in the accident area not more than  $\pm 1$  unit,

state  $r_{(1)}^{12}$  – the realised substance changed pH regime in the accident area not more than  $\pm 1$  unit, and simultaneously the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}/2, LC_{50}>$ ,

state  $r_{(1)}^{13}$  – the realised substance changed pH regime in the accident area not more than  $\pm 1$  unit, and simultaneously the realised substance caused the aesthetic nuisance of air of the accident area but the closure of area is not required, and additionally the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(0, LC_{50}/2>$ ,

state  $r_{(1)}^{14}$  – the realised substance changed pH regime in the accident area  $\pm 1-2$  units, and simultaneously the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for not more than 2 days, and additionally the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}/2, LC_{50}>$ ,

state  $r_{(1)}^{15}$  – the realised substance changed pH regime in the accident area  $\pm 1-2$  units, and simultaneously the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for not more than 2 days, and additionally the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}, \infty)$ ,

state  $r_{(1)}^{16}$  – the realised substance changed pH regime in the accident area  $\pm 1-2$  units, and simultaneously the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for 2 days or more, and additionally the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}, \infty)$ ,

state  $r_{(1)}^{17}$  – the realised substance decreased oxygen concentration in the air of the accident area of the value up to 2%,

state  $r_{(1)}^{18}$  – the realised substance decreased oxygen concentration in the air of the accident area of the value from the interval 2-5%,

state  $r_{(1)}^{19}$  – the realised substance decreased oxygen concentration in the air of the accident area of the value more than 5%,

state  $r_{(1)}^{20}$  – the realised substance increased the air temperature in the accident area of the value from the interval  $(10^{\circ}\text{C}, 20^{\circ}\text{C}>$ ,

state  $r_{(1)}^{21}$  – the realised substance increased the air temperature in the accident area of the value from the interval  $(10^{\circ}\text{C}, 20^{\circ}\text{C}>$ , and simultaneously the realised substance decreased oxygen concentration in the air of the accident area of the value up to 2%, and additionally the realised substance caused the

aesthetic nuisance of air of the accident area but the closure of area is not required,

state  $r_{(1)}^{22}$  – the realised substance increased the air temperature in the accident area of the value from the interval  $(10^{\circ}\text{C}, 20^{\circ}\text{C}>$ , and simultaneously the realised substance decreased oxygen concentration in the air of the accident area of the value up to 2%, and additionally the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for not more than 2 days, and moreover the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}/2, LC_{50}>$ ,

state  $r_{(1)}^{23}$  – the realised substance increased the air temperature in the accident area of the value from the interval  $(20^{\circ}\text{C}, 30^{\circ}\text{C}>$ , and simultaneously the realised substance decreased oxygen concentration in the air of the accident area of the value up to 2%,

state  $r_{(1)}^{24}$  – the realised substance increased the air temperature in the accident area of the value from the interval  $(20^{\circ}\text{C}, 30^{\circ}\text{C}>$ , and simultaneously the realised substance decreased oxygen concentration in the air of the accident area of the value from the interval 2-5% and additionally the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for not more than 2 days,

state  $r_{(1)}^{25}$  – the realised substance increased the air temperature in the accident area of the value from the interval  $(20^{\circ}\text{C}, 30^{\circ}\text{C}>$ , and simultaneously the realised substance decreased oxygen concentration in the air of the accident area of the value from the interval 2-5%, and additionally the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for 2 days or more, and moreover the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}, \infty)$ ,

state  $r_{(1)}^{26}$  – the realised substance increased the air temperature in the accident area of the value more than  $30^{\circ}\text{C}$ , and simultaneously the realised substance decreased oxygen concentration in the air of the accident area of the value from the interval 2-5%,

state  $r_{(1)}^{27}$  – the realised substance increased the air temperature in the accident area of the value more than  $30^{\circ}\text{C}$ , and simultaneously the realised substance decreased oxygen concentration in the air of the accident area of the value more than 5%, and additionally the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for 2 days or more,

state  $r_{(1)}^{28}$  – the realised substance increased the air temperature in the accident area of the value more than 30°C, and simultaneously the realised substance decreased oxygen concentration in the air of the accident area of the value more than 5%, and additionally the realised substance caused the aesthetic nuisance of air of the accident area and the closure of area is required for 2 days or more, and moreover the realised substance caused the pollution and its concentration in the air in the accident area belongs to the interval  $(LC_{50}, \infty)$ ,

state  $r_{(1)}^{29}$  – the realised substance increased the air temperature in the accident area of the value from the interval  $(20^\circ\text{C}, 30^\circ\text{C})$ ,

state  $r_{(1)}^{30}$  – the realised substance increased the air temperature in the accident area of the value more than 30°C, and simultaneously the realised substance decreased oxygen concentration in the air of the accident area of the value up to 2%.

Moreover, there are  $\ell_2 = 28$  states for the water surface ( $D_2$  sub-region),  $\ell_3 = 28$  states for the water column ( $D_3$  sub-region),  $\ell_4 = 31$  states for the sea floor ( $D_4$  sub-region), and  $\ell_5 = 23$  states for the coast ( $D_5$  sub-region) that are given in [EU-CIRCLE Report D3.3-GMU21, 2016].

### 2.1.2. Probabilities of transitions between states of the process of environment degradation

On the basis of the statistical data, it is possible to evaluate the following unknown basic parameters of the process of environment degradation at the Baltic Sea waters:

- the vectors of the initial probabilities  $q_{(k/v)}^i(0)$  of the environment degradation sub-process at the particular states at the moment  $t = 0$  as follows:

$$[q_{(1/v)}(0)]_{1 \times 30} = [1, 0, \dots, 0]$$

for  $v = 1, 6, 27, 30$ ,

$$[q_{(2/v)}(0)]_{1 \times 28} = [1, 0, \dots, 0]$$

for  $v = 1, 17, 33$ ,

$$[q_{(3/v)}(0)]_{1 \times 28} = [1, 0, \dots, 0]$$

for  $v = 1, 14, 24$ ,

$$[q_{(4/v)}(0)]_{1 \times 31} = [1, 0, \dots, 0]$$

for  $v = 1, 14$ ,

$$[q_{(5/v)}(0)]_{1 \times 23} = [1, 0, \dots, 0]$$

for  $v = 1$ ,

and

$$[q_{(1/v)}(0)]_{1 \times 30} = [0, 0, \dots, 0]$$

for  $v = 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 28, 29, 31, 32, 33, 34, 35$ ,

$$[q_{(2/v)}(0)]_{1 \times 28} = [0, 0, \dots, 0]$$

for  $v = 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32$ ,

$$[q_{(3/v)}(0)]_{1 \times 28} = [0, 0, \dots, 0]$$

for  $v = 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28, 29$ ,

$$[q_{(4/v)}(0)]_{1 \times 31} = [0, 0, \dots, 0],$$

for  $v = 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29$ ,

$$[q_{(5/v)}(0)]_{1 \times 23} = [0, 0, \dots, 0],$$

for  $v = 2, 3, \dots, 29$ ; (1)

- the matrix  $[q_{(k/v)}^{ij}]$   $i, j = 1, 2, \dots, \ell_k$ ,  $k = 1, 2, \dots, 5$ ,  $v = 1, 2, \dots, v_k$ , of the probabilities of transitions of the sub-process  $R_{(k/v)}(t)$  transitions from the state  $r_{(k/v)}^i$ , into the state  $r_{(k/v)}^j$ , during the experimental time. The probabilities of transitions that are not equal to 0 are as follows:

$$q_{(1/6)}^{12} = 1, q_{(1/6)}^{21} = 1; \quad (2)$$

$$q_{(1/27)}^{16} = 1, q_{(1/27)}^{61} = 1; \quad (3)$$

$$q_{(1/30)}^{111} = 1, q_{(1/30)}^{111} = 1; \quad (4)$$

$$q_{(2/17)}^{1\ 27} = 1, q_{(2/17)}^{12\ 1} = 1, q_{(2/17)}^{16\ 12} = 1, q_{(2/17)}^{21\ 16} = 1, \\ q_{(2/17)}^{25\ 21} = 1, q_{(2/17)}^{27\ 25} = 1; \quad (5)$$

$$q_{(2/33)}^{1\ 6} = 1, q_{(2/33)}^{6\ 1} = 1; \quad (6)$$

$$q_{(3/14)}^{1\ 27} = 1, q_{(3/14)}^{12\ 1} = 1, q_{(3/14)}^{16\ 12} = 1, q_{(3/14)}^{21\ 16} = 1, \\ q_{(3/14)}^{25\ 21} = 1, q_{(3/14)}^{27\ 25} = 1; \quad (7)$$

$$q_{(3/24)}^{1\ 6} = 1, q_{(3/24)}^{6\ 1} = 1; \quad (8)$$

$$q_{(4/14)}^{1\ 30} = 1, q_{(4/14)}^{12\ 1} = 1, q_{(4/14)}^{16\ 12} = 1, q_{(4/14)}^{21\ 16} = 1, \\ q_{(4/14)}^{28\ 21} = 1, q_{(4/14)}^{30\ 28} = 1. \quad (9)$$

The values of some probabilities existing in the vectors  $[q_{(k/v)}(0)]$  and in the matrix  $[q_{(k/v)}^{ij}]$ , besides of those standing on the main diagonal, equal to zero do not mean that the events they are concerned with, cannot appear. They evaluated on the basis of real statistical data and their values may change and become more precise if the duration of the experiment is longer.

### 2.1.3. Evaluation of distributions and mean values of the process of environment degradation conditional sojourn times

Because we only have the number of realizations of the sub-process of environment degradation and its all realizations are equal to an approximate value, we assume that this time has the uniform distribution in the interval from this value minus its half to this value plus its half.

The uniform distribution functions of the sub-process of environment degradation for particular conditional sojourn times  $\zeta_{(k/v)}^{ij}$  are identified on the basis of statistical data coming from its process realizations at the Baltic Sea waters given in Appendix 6 in [EU-CIRCLE Report D3.3-GMU22, 2016]. For instance, the sub-process of environment degradation the conditional sojourn time  $\zeta_{(1/6)}^{12}$  assumed  $n_{(1/6)}^{12} = 1$  value equals to 1, we assume that it has the uniform distribution function given by

$$G_{(1/6)}^{12}(t) = \begin{cases} 0, & t < 0.5 \\ t, & 0.5 \leq t < 1.5 \\ 1, & t \geq 1.5. \end{cases} \quad (10)$$

In the case when as a result of the experiment, coming from experts, we have less than 28 realizations of the

process of environment degradation, we determined this conditional sojourn times have the empirical distributions. The empirical distribution functions of the process of environment degradation for particular conditional sojourn times  $\zeta_{(k/v)}^{ij}$  are identified on the basis of statistical data coming from its process realizations at the Baltic Sea waters given in Appendix 6 in [EU-CIRCLE Report D3.3-GMU22, 2016]. For instance, the process initiating events conditional time  $\zeta_{(1/27)}^{61}$  assumed  $n_{(1/27)}^{61} = 2$  values. The order sample realizations  $\zeta_{(1/27)}^{61}$  is: 180, 300. Thus, we assume that conditional sojourn time  $\zeta_{(1/27)}^{61}$  has the empirical distribution function given by

$$G_{(1/27)}^{61}(t) = \begin{cases} 0, & t < 180 \\ 1/2, & 180 \leq t < 300 \\ 1, & t \geq 300. \end{cases} \quad (11)$$

We have proceeded with the remaining conditional times at the states of the sub-process of environment degradation in the same way and approximately fix they distribution.

Further, for distributions identified in this section the approximate empirical values of the mean values  $M_{(k/v)}^{ij} = E[\zeta_{(k/v)}^{ij}]$ ,  $i, j = 1, 2, \dots, \ell_k$ ,  $i \neq j$ ,  $k = 1, 2, \dots, 5$ ,  $\ell_1 = 30$ ,  $\ell_2 = 28$ ,  $\ell_3 = 28$ ,  $\ell_4 = 31$ ,  $\ell_5 = 23$ ,  $\nu_1 = 35$ ,  $\nu_2 = 33$ ,  $\nu_3 = 29$ ,  $\nu_4 = 29$ ,  $\nu_5 = 29$ , of the process of environment degradation conditional sojourn times at particular states at the Baltic Sea waters can be determined and they are as follows:

$$M_{(1/6)}^{1\ 2} = 1, M_{(1/6)}^{2\ 1} = 240, M_{(1/27)}^{1\ 6} = 1, M_{(1/27)}^{6\ 1} = 240,$$

$$M_{(1/30)}^{1\ 11} = 1, M_{(1/30)}^{11\ 1} = 240, M_{(2/17)}^{1\ 27} = 1,$$

$$M_{(2/17)}^{12\ 1} = 2880, M_{(2/17)}^{16\ 12} = 3780, M_{(2/17)}^{21\ 16} = 2880,$$

$$M_{(2/17)}^{25\ 21} = 300, M_{(2/17)}^{27\ 25} = 240, M_{(2/33)}^{1\ 6} = 1,$$

$$M_{(2/33)}^{6\ 1} = 1440, M_{(3/14)}^{1\ 27} = 1, M_{(3/14)}^{12\ 1} = 2880,$$

$$M_{(3/14)}^{16\ 12} = 3780, M_{(3/14)}^{21\ 16} = 2880, M_{(3/14)}^{25\ 21} = 300,$$

$$M_{(3/14)}^{27\ 25} = 240, M_{(3/24)}^{1\ 6} = 1, M_{(3/24)}^{6\ 1} = 1440,$$

$$M_{(4/14)}^{1\ 30} = 1, M_{(4/14)}^{12\ 1} = 2880, M_{(4/14)}^{16\ 12} = 3780,$$

$$M_{(4/14)}^{21\ 16} = 2880, M_{(4/14)}^{28\ 21} = 300, M_{(4/14)}^{30\ 28} = 240. \quad (12)$$

#### 2.1.4. Prediction of the process of environment degradation

Using the identified parameters of the process of environment degradation in Section 2.1.2 and 2.1.3, it is possible to predict its characteristics [EU-CIRCLE Report D3.3-GMU23, 2016]. Namely, considering (2)-(9) and (12), the mean values of the process of environment degradation at the Baltic Sea waters unconditional sojourn times at the particular states are:

$$M_{(1/6)}^1 = 1, M_{(1/6)}^2 = 240; \quad (13)$$

$$M_{(1/27)}^1 = 1, M_{(1/27)}^6 = 240; \quad (14)$$

$$M_{(1/30)}^1 = 1, M_{(1/30)}^{11} = 240; \quad (15)$$

$$M_{(2/17)}^1 = 1, M_{(2/17)}^{12} = 2880, M_{(2/17)}^{16} = 3780, \\ M_{(2/17)}^{21} = 2880, M_{(2/17)}^{25} = 300, M_{(2/17)}^{27} = 240; \quad (16)$$

$$M_{(2/33)}^1 = 1, M_{(2/33)}^6 = 1440; \quad (17)$$

$$M_{(3/14)}^1 = 1, M_{(3/14)}^{12} = 2880, M_{(3/14)}^{16} = 3780, \\ M_{(3/14)}^{21} = 2880, M_{(3/14)}^{25} = 300, M_{(3/14)}^{27} = 240; \quad (18)$$

$$M_{(3/24)}^1 = 1, M_{(3/24)}^6 = 1440; \quad (19)$$

$$M_{(4/14)}^1 = 1, M_{(4/14)}^{12} = 2880, M_{(4/14)}^{16} = 3780, \\ M_{(4/14)}^{21} = 2880, M_{(4/14)}^{28} = 300, M_{(4/14)}^{30} = 240. \quad (20)$$

Since from the system of equations (4.30) in [EU-CIRCLE Report D3.3-GMU21, 2016] takes the following form

$$\begin{cases} [\pi_{(k/v)}^i]_{1 \times \ell_k} = [\pi_{(k/v)}^i]_{1 \times \ell_k} [q_{(k/v)}^{ij}]_{\ell_k \times \ell_k} \\ \sum_{j=1}^{\ell_k} \pi_{(k/v)}^j = 1, \end{cases}$$

where

$$i, j = 1, 2, \dots, \ell_k, i \neq j, k = 1, 2, \dots, 5, \ell_1 = 30, \ell_2 = 28, \\ \ell_3 = 28, \ell_4 = 31, \ell_5 = 23, \nu_1 = 35, \nu_2 = 33, \nu_3 = 29, \\ \nu_4 = 29, \nu_5 = 29,$$

we get its following solution:

$$\pi_{(1/6)}^1 \cong 0.5, \pi_{(1/6)}^2 \cong 0.5; \quad (21)$$

$$\pi_{(1/27)}^1 = 0.5, \pi_{(1/27)}^6 = 0.5; \quad (22)$$

$$\pi_{(1/30)}^1 = 0.5, \pi_{(1/30)}^{11} = 0.5; \quad (23)$$

$$\pi_{(2/17)}^1 \cong 0.1667, \pi_{(2/17)}^{12} \cong 0.1667, \pi_{(2/17)}^{16} \cong 0.1667, \\ \pi_{(2/17)}^{21} \cong 0.1667, \pi_{(2/17)}^{25} \cong 0.1666, \\ \pi_{(2/17)}^{27} \cong 0.1666; \quad (24)$$

$$\pi_{(2/33)}^1 \cong 0.5, \pi_{(2/33)}^6 \cong 0.5; \quad (25)$$

$$\pi_{(3/14)}^1 \cong 0.1667, \pi_{(3/14)}^{12} \cong 0.1667, \pi_{(3/14)}^{16} \cong 0.1667, \\ \pi_{(3/14)}^{21} \cong 0.1667, \pi_{(3/14)}^{25} \cong 0.1666, \\ \pi_{(3/14)}^{27} \cong 0.1666; \quad (26)$$

$$\pi_{(3/24)}^1 \cong 0.5, \pi_{(3/24)}^2 \cong 0.5; \quad (27)$$

$$\pi_{(4/14)}^1 \cong 0.1667, \pi_{(4/14)}^{12} \cong 0.1667, \pi_{(4/14)}^{16} \cong 0.1667, \\ \pi_{(4/14)}^{21} \cong 0.1667, \pi_{(4/14)}^{28} \cong 0.1666, \\ \pi_{(4/14)}^{30} \cong 0.1666. \quad (28)$$

Then after considering (13)-(20) respectively and applying (4.29) in [EU-CIRCLE Report D3.3-GMU21, 2016] we get the approximate limit values of transient probabilities at the particular states of the process of environment degradation:

$$q_{(1/6)}^1 = 0.00415, q_{(1/6)}^2 = 0.99585; \quad (29)$$

$$q_{(1/27)}^1 = 0.00415, q_{(1/27)}^6 = 0.99585; \quad (30)$$

$$q_{(1/30)}^1 = 0.00415, q_{(1/30)}^{11} = 0.99585; \quad (31)$$

$$q_{(2/17)}^1 = 0.00010, q_{(2/17)}^{12} = 0.28570, \\ q_{(2/17)}^{16} = 0.37497, q_{(2/17)}^{21} = 0.28570, \\ q_{(2/17)}^{25} = 0.02974, q_{(2/17)}^{27} = 0.02379; \quad (32)$$

$$q_{(2/33)}^1 = 0.00069, q_{(2/33)}^6 = 0.99931; \quad (33)$$

$$\begin{aligned} q_{(3/14)}^1 &= 0.00010, & q_{(3/14)}^{12} &= 0.28570, \\ q_{(3/14)}^{16} &= 0.37497, & q_{(3/14)}^{21} &= 0.28570, \\ q_{(3/14)}^{25} &= 0.02974, & q_{(3/14)}^{27} &= 0.02379; \end{aligned} \quad (34)$$

$$q_{(3/24)}^1 = 0.00069, \quad q_{(3/24)}^6 = 0.99931; \quad (35)$$

$$\begin{aligned} q_{(4/14)}^1 &= 0.00010, & q_{(4/14)}^{12} &= 0.28570, \\ q_{(4/14)}^{16} &= 0.37497, & q_{(4/14)}^{21} &= 0.28570, \\ q_{(4/14)}^{28} &= 0.02974, & q_{(4/14)}^{30} &= 0.02379. \end{aligned} \quad (36)$$

Further, by (4.31) in [EU-CIRCLE Report D3.3-GMU21, 2016] and considering (29)-(36) respectively, the approximate mean values of the sojourn total times  $\hat{\zeta}^i$  of the sub-processes of environment degradation  $R_{(k/v)}(t)$  in the time interval  $\zeta = 1 \text{ month} = 43200 \text{ minutes}$  at the particular states  $r_{(k/v)}^i$ , expressed in minutes are:

$$\hat{M}_{(1/6)}^1 = 179.25, \quad \hat{M}_{(1/6)}^2 = 43020.75; \quad (37)$$

$$\hat{M}_{(1/27)}^1 = 179.25, \quad \hat{M}_{(1/27)}^6 = 43020.75; \quad (38)$$

$$\hat{M}_{(1/30)}^1 = 179.25, \quad \hat{M}_{(1/30)}^{11} = 43020.75; \quad (39)$$

$$\begin{aligned} \hat{M}_{(2/17)}^1 &= 4.29, & \hat{M}_{(2/17)}^{12} &= 12342.03, \\ \hat{M}_{(2/17)}^{16} &= 16198.91, & \hat{M}_{(2/17)}^{21} &= 12342.03, \\ \hat{M}_{(2/17)}^{25} &= 1284.86, & \hat{M}_{(2/17)}^{27} &= 1027.89; \end{aligned} \quad (40)$$

$$\hat{M}_{(2/33)}^1 = 29.98, \quad \hat{M}_{(2/33)}^6 = 43170.02; \quad (41)$$

$$\begin{aligned} \hat{M}_{(3/14)}^1 &= 4.29, & \hat{M}_{(3/14)}^{12} &= 12342.03, \\ \hat{M}_{(3/14)}^{16} &= 16198.91, & \hat{M}_{(3/14)}^{21} &= 12342.03, \\ \hat{M}_{(3/14)}^{25} &= 1284.86, & \hat{M}_{(3/14)}^{27} &= 1027.89; \end{aligned} \quad (42)$$

$$\hat{M}_{(3/24)}^1 = 29.98, \quad \hat{M}_{(3/24)}^6 = 43170.02; \quad (43)$$

$$\begin{aligned} \hat{M}_{(4/14)}^1 &= 4.29, & \hat{M}_{(4/14)}^{12} &= 12342.03, \\ \hat{M}_{(4/14)}^{16} &= 16198.91, & \hat{M}_{(4/14)}^{21} &= 12342.03, \\ \hat{M}_{(4/14)}^{28} &= 1284.86, & \hat{M}_{(4/14)}^{30} &= 1027.89. \end{aligned} \quad (44)$$

### 3. Conclusion

The results (29)-(36) and (37)-(44) are main characteristics of the considered process of environment degradation that is the third part of the integrated model of critical infrastructure accident consequences [Bogalecka, Kołowrocki, 2017c]. This characteristics together with results obtained in [Bogalecka, Kołowrocki, 2017a,b] can be used to the prediction of critical infrastructure accident losses [Bogalecka, Kołowrocki, 2017d].

### Acknowledgments



The paper presents the results developed in the scope of the EU-CIRCLE project titled “A pan – European framework for strengthening Critical Infrastructure resilience to climate change” that has received funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement No 653824. <http://www.eu-circle.eu/>.

### References

- Bogalecka M., Analysis of sea accidents initial events, Polish Journal of Environmental Studies, 19(4A), 5-8, 2010
- Bogalecka M., Kołowrocki K., Modelling, identification and prediction of environment degradation initial events process generated by critical infrastructure accidents. Journal of Polish Safety and Reliability Association, Summer Safety and Reliability Seminars, 6(1), 47-66, 2015a
- Bogalecka M., Kołowrocki K., The process of sea environment threats generated by hazardous chemicals release. Journal of Polish Safety and Reliability Association, Summer Safety and Reliability Seminars, 6(1), 67-74, 2015b
- Bogalecka M., Kołowrocki K., Modelling critical infrastructure accident consequences – an overall approach. Journal of Polish Safety and Reliability Association, Summer Safety and Reliability Seminars, 7(1), 1-13, 2016
- Bogalecka M., Kołowrocki K., General model of critical infrastructure accident consequences application to chemical spill consequences generated by dynamic ship critical infrastructure network operating at the Baltic Sea waters. Part 1. Process of initiating events. Journal of Polish Safety and Reliability Association, Summer Safety and Reliability Seminars, 8(3), 117-121, 2017a
- Bogalecka M., Kołowrocki K., General model of critical infrastructure accident consequences

application to chemical spill consequences generated by dynamic ship critical infrastructure network operating at the Baltic Sea waters. Part 2. Process of environment threats. Journal of Polish Safety and Reliability Association, Summer Safety and Reliability Seminars, 8(3), 123-129, 2017b

Bogalecka M., Kołowrocki K., Integrated model of critical infrastructure accident consequences. Journal of Polish Safety and Reliability Association, Summer Safety and Reliability Seminars, 8(3), 43-52, 2017c

Bogalecka M., Kołowrocki K., Integrated impact model on critical infrastructure accident consequences related to climate-weather change process. Journal of Polish Safety and Reliability Association, Summer Safety and Reliability Seminars, 8(4), 67-75, 2017d

EU-CIRCLE Report D3.3-GMU21, Modelling critical infrastructure accident consequences – designing the General Model of Critical Infrastructure Accident Consequences (GMCIAC), 2016

EU-CIRCLE Report D3.3-GMU22, Identification of unknown parameters of the General Model of Critical Infrastructure Accident Consequences (GMCIAC), 2016

EU-CIRCLE Report D3.3-GMU23, Adaptation of the general model of critical infrastructure accident consequences (GMCIAC) to the prediction of critical infrastructure accident consequences, 2016

Grabski F., Semi-Markov processes: applications in system reliability and maintenance. Elsevier, 2015

Kołowrocki K., Reliability of large systems. Amsterdam, Boston, Heidelberg, London, New York, Oxford, Paris, San Diego, San Francisco, Singapore, Sidney, Tokyo, Elsevier, 2004

Kołowrocki K., Reliability of large and complex systems. Amsterdam, Boston, Heidelberg, London, New York, Oxford, Paris, San Diego, San Francisco, Singapore, Sidney, Tokyo, Elsevier, 2014

Kołowrocki K., Soszyńska-Budny J., A general model of industrial systems operation processes related to their environment and infrastructure. Journal of Polish Safety and Reliability Association, Summer Safety and Reliability Seminars, 2(2), 223-226, 2008

Kołowrocki K., Soszyńska-Budny J., Reliability and safety of complex technical systems and processes: modeling – identification – prediction – optimization. London, Dordrecht, Heildeberg, New York, Springer, 2011

Limnios N., Oprisan G., Semi-Markov processes and reliability. Birkhauser, Boston, 2005

Macci C., Large deviations for empirical estimators of the stationary distribution of a semi-Markov process with finite state space. Communications in Statistics-Theory and Methods, 37(9), 3077-3089, 2008

Mercier S., Numerical bounds for semi-Markovian quantities and application to reliability. Methodology and Computing in Applied Probability, 10(2), 179-198, 2008